



Suggested Climate Change Adaptation Strategies for The Narrows WMA, Stands 1 and 5

Andrew Whitman

Introduction

Climate change will alter many aspects of forests and forest habitats in Vermont (Frumhoff et al. 2007). Although land managers cannot control changes in climate (e.g., warmer temperatures, altered precipitation), they have many options for increasing the resiliency of forests (Gunn et al. 2009). Managing forests under climate change is a challenging undertaking, as the timing and severity of anticipated changes are hard to predict and vary regionally. Climate change adaptation is the planning and application of policies and practices that reduce the threats and impacts posed by climate change. The goal of climate change adaptation in working forests is not to stop climate impacts or preserve the current forest composition, but rather to maintain key forest values. The overarching goals of climate change adaptation in The Narrows WMA are to maintain forest cover and its many economic, social, and ecological values, and achieve the goals established for this management unit (Appendix A).

Summary of Projected Climate Change Impacts on Southern Vermont Forests

Deciduous and mixed forests in Vermont may change significantly in the next 100 years under every climate change scenario (Rustad et al. 2012, Prasad et al. 2007). The extent of oak and pine forest types (including oak-hickory types) is projected to increase and further expand in southern Vermont (Iverson et al. 2008a). Under the lowest emissions scenario, Vermont is predicted to retain its northern hardwood forests. Northern hardwood tree species may achieve increased growth rates under low- and moderate-emissions scenarios due to higher temperatures and a longer growing season, potential CO₂-driven increases in photosynthesis and water-use efficiency, and changes in




Figure 1. The Narrows Wildlife Management Area and approximate location of the proposed harvest area (shaded).

(N) cycle that increase N availability and plant productivity (Butler et al. 2012). If CO₂ fertilization does not occur, growth rates are projected to increase slightly. Under the higher emissions scenario, growth rates of northern hardwood tree species may decline by 2100 due to temperature stress (Ollinger et al. 2008). Under high-emissions scenarios, oak-hickory forest types are also

the
nitrogen

projected to increase greatly in southern Vermont and northern hardwood forest will modestly decline (Tang and Beckage 2010). Hemlock woolly adelgid (*Adelges tsugae*) is projected to expand into the entire southern Vermont area in the next 30 years and could eliminate eastern hemlock (*Tsuga canadensis*) (Paradis et al. 2008). Several northern tree species, including sugar maple (*Acer saccharum*), yellow birch (*Betula alleghaniensis*), and ash spp. (*Fraxinus* spp.), have had periods of decline or reduced productivity in the past 100 years (Mohan et al. 2009).

It is unclear how forest stands and plant and animal communities will respond to climate change. Although there is the potential for large climate-driven range shifts in forest species and types by 2100 (Iverson et al. 2008a), species shifts are not expected to keep up with the rate of climate change, and will likely be delayed (Mohan et al. 2009). Competition may be the dominant structuring force for tree communities in forests (Zhu et al. 2012), so climate change impacts may have to be large to result in changes in regional distributions of tree species. Some tree species may take >100 years before they begin to colonize significant portions of new habitat (Iverson et al. 2005). Therefore, many present and future forest communities may be composed of plant species with migration rates far below those required to track contemporary climate change (Zhu et al. 2012).

Extreme weather (e.g., high temperatures and ozone levels, drought, late frosts, ice storms) might increase mortality that varies depending on position on slope, site index, soils, and age classes (trees, saplings, and seedlings). For many species on mesic sites, it may be a slow replacement of seedlings and saplings of northern species with those of southern species. For some highly vulnerable species (e.g., eastern hemlock), catastrophic mortality may greatly reduce overstory and possible lack of regeneration could make it difficult to restore mature forest conditions in the long term.

Proposed Overall Management Approach

Stand Number	Stand Type	Silvicultural Approach	Wildlife Approach ¹	Composition ¹	Cycle of Entries
5	oak/hardwood	uneven-aged management	Maintain mast component and natural community composition.	w. pine = 19% s. maple = 26% red oak = 7% s. hickory = <1%	15-year cutting cycle
1	white pine	uneven-aged management	Maintain softwood cover but do not rely on hemlock for future cover.	w. pine = 83%	15-year to 25-year cutting cycle

¹ The Narrows Wildlife Management Area: Long Range Management Plan (Maciejowski et al 2011).

Projected Climate Change Impacts

Temperature

Temperatures are projected to increase and reduce the productivity of northern hardwood species late in the century. This is expected to slightly increase mortality of overstory trees and increase their susceptibility to pests and disease (Hanson and Weltzin 2000). Southern species (e.g., shagbark hickory, oak spp.) are more resistant to high temperatures (Burns and Honkala 1990). Seedling mortality of northern conifers (spruce, fir, cedar) is greatest at temperatures >90° F (Burns and Honkala 1990). The expected impact of future temperature increases on this site is moderate and its likelihood is high.

Drought

The frequency of short-term (one to three months) drought is projected to at least double (Hayhoe et al. 2007). Drought is expected to increase mortality of overstory trees and increase their susceptibility to pests and disease (Allen et al 2010, Hanson and Weltzin 2000). The well-drained nature of the local soils will likely make vegetation susceptible to drought, except where sub-surface moisture collects in seeps. The expected impact of future drought on this site is moderate and its likelihood is high.

Lack of snow cover

In northern hardwood and spruce-fir forests, the impact of droughts could increase greatly when preceded by winters during which snow cover is inadequate (<25 cm) and intense cold spells occur, making root systems more vulnerable to root kill (Auclair et al. 2010). The expected impact of future lack of snow cover on this site is low to moderate and its likelihood is high.

Wildfire

Climate change may increase the frequency of lightning strikes, dryness, and blocking high pressure ridges, which rapidly dry forest fuels, thereby increasing the probability and extent of fire in Vermont. Modeled predictions suggest

that fire severity due to changing weather conditions may increase 10% to 20% overall for the Northeast (Flannigan et

al. 2009). Under a range of scenarios, the probability and extent of hardwood forest burned in nearby Quebec was projected to range from a slight increase to a five-fold increase (Drever et al. 2009). Fire in hardwood forests could increase if conditions lead to long, warm periods before leaf out or after leaf fall. The expected impact of future wildfire on this site is moderate to high and its likelihood is low.

Current Challenges

Lack of natural regeneration

Natural regeneration of tree seedlings and saplings in Stands 1 and 5 were sparse and were heavily browsed. Climate change by itself may not significantly affect growth and survival of seedlings and saplings of most species, except for spruce, northern white cedar, and fir seedlings, which can have high mortality in hot and dry years (Burns and Honkala 1990).

Exotic plant species

Invasive exotic plant species were common in Stands 1 and 5 and their presence is a major management challenge. Several exotic species are well established and have significant impact, including Japanese honeysuckle, buckthorn, and barberry. These species aggressively increased after harvest on previously harvested sites. Climate change is predicted to increase the frequency of disturbance events that lead to rapid expansion of exotic species and changes in plant communities (Dale et al. 2001). Because most exotic plant species have high growth rates and long-distance dispersal traits, they have a competitive advantage over native species for colonizing and establishment, especially following ecosystem disturbance (Dukes and Mooney 1999). Climate will hasten the spread of many exotic plant species because (1) increased forest ecosystem disturbance and stress will facilitate their establishment and (2) rising CO₂ levels enhance the growth and competitiveness of some exotic species (Ziska and Dukes 2011).

Hemlock Woolly Adelgid (HWA)

HWA is not present at The Narrows WMA, but is found in southern Vermont (VT Department of Forests, Parks, and Recreation 2012). With climate change, winters are projected to become milder and so HWA is predicted to occupy The Narrows WMA in the near future (Dukes et al. 2009).

Deer

High deer population densities have resulted in moderate to heavy browsing of all understory species, except exotic

species, in the unit (Maciejowski et al. 2011), including Stands 1 and 5. As a result, the shrub layer, including tree regeneration, is largely absent in many areas. Deer also avoid browsing on exotic plant species to the detriment of native species (Knight et al. 2009). With climate change, winter will be milder and deer population densities can be expected to increase (Rustad et al. 2012).

Soils

The soils of the harvest area are silt loams that are part of the Farmington-Galway-Galoo complex (USDA NRCS 2013). They are 10 to 20 inches deep to bedrock, somewhat excessively drained, and have very low available water capacity and so these stands are somewhat susceptible to drought. Short-term drought may increase significantly with climate change but may only affect growth and regeneration survival. Severe droughts (2-5 years in length) may weaken canopy trees and lead to their mortality but will only increase slightly with climate change. Northern conifers (spruce, northern white cedar, and fir), Eastern hemlock, and yellow birch are sensitive to drought and their regeneration may be vulnerable to short-term drought on shallow soils (Burns and Honkala 1990). Maple spp. are somewhat drought tolerant. Species that most drought resistant on these sites include white pine, shagbark hickory, and red oak (Burns and Honkala 1990).

Specific Suggestions for the Harvest Plan

Climate change is expected to increase air temperatures, frequency of drought and forest stress, and pest outbreaks. With these changes, invasive species, tree diseases, and tree pests are projected to become more widespread and threaten forests. Storm damage is projected to increase as well. Therefore, for this site, climate change adaptation focuses on using standard forest practices to achieve greater stand resiliency and vigor and reduce vulnerability. The proposed approach focuses on Management Goals 1 through 3, 6, and 8 (Appendix A).

Almost all of these goals and practices are either already included in the existing plan for The Narrows or are consistent with its goals. Climate change adaptation often simply requires finding opportunities to make small changes to existing landowner goals and practices that increase stand resilience to climate change

Long term Strategies

This section identifies how the existing plan has goals that are consistent with climate change adaptation.

1. Increase stand diversity and health

Stand resiliency can be achieved by maintaining a diversity of tree ages and species in the stand. This reduces stand vulnerability to losses of age classes and species that are susceptible to climate change-related stressors.

a. Aim for diverse-age structure: The stands should be shifted to an uneven-age structure to allow advanced regeneration for in fill if small or large replacement events occur and improve flexibility for managing the stand for financial and biodiversity goals in the future. A diversity of ages means that the stand always has younger, vigorously growing stock (Wilkerson et al. 2011, Gunn et al. 2009). The current plan also suggests this strategy for these stands.

b. Maintain tree species diversity: Tree species diversity at the stand level should be maintained to improve stand-level disease resistance and create greater flexibility for managing stands for a range of silvicultural, financial, and biodiversity goals in the future (Wilkerson et al. 2011). This includes maintaining shagbark hickory in the stand, a southern species that has been projected to increase its range in Vermont and that should be more resilient to future climate conditions than northern species. It also means reducing eastern hemlock densities where it dominates to allow other tree species to enter the canopy and provide canopy cover if HWA kills many eastern hemlock trees. So for Stand 1, where the natural community would be hemlock, aim for a hemlock-northern hardwood mix as a resilience strategy. This goal is consistent with goals for the WMA, including these stands.

2. Reduce the impact of existing stressors on forests

Reducing existing stressors can make ecosystems less vulnerable to stress posed by climate change.

Control invasive understory species: Invasive understory species should be controlled to allow recruitment of regeneration and for the stand in the future (Wilkerson et al. 2011). Control should target portions of the harvest block where harvesting will create canopy gaps and should be applied before and, if necessary, after the harvest (VT Fish and Wildlife Department 2012). Control can include use of backpack sprayers or manual removal to target large and/or vigorous invasive shrubs. This goal is part of the current plan for the WMA, including these stands.

3. Protect soils

Protecting soils is essential for maintaining site productivity. Climate change will likely increase the frequency of extreme events and necessitate consistent use of measures to protect soils from erosion and rutting.

Maintain adequate down woody materials: Developing or maintaining adequate down woody material will replenish soil organic matter, moderate temperatures, and support wildlife habitat. Productive and structurally diverse stands will better adapt to unknown site changes likely to result from climate change. Tops and slash should be placed around stumps of oak and maple species to protect sprouts from deer browsing. This goal is part of the current plan for the WMA, including these stands.

4. Reduce impacts of current stressors to wildlife

By reducing the threats posed by existing stressors, one can reduce the vulnerability of wildlife to climate change. This includes:

- Control the spread of invasive exotic species.
- Maintain/enhance mast component by harvesting to promote crown development of species including oaks and hickories.
- Maintain or enhance the mosaic of forest stands and natural communities for their contribution to wildlife habitat including softwood cover and mast source.
- Protect vernal pools and their habitat value by following vernal pool management guidelines.
- Protect seeps and intermittent drainages where plant diversity is high and uncommon plant species occur more frequently.

Short-term Silvicultural and Operational Practices

All of these strategies were discussed on the ground with the Stewardship Team or management staff and were consistent with existing discussions and strategies for management. Most of these practices have been routinely used on state lands.

1. Apply crop tree release (addresses Goals 6 and 8)

Crop tree release should be applied to grow very large trees in the future for wildlife habitat (e.g., bat roosts, habitat for mature forest species); maintain a vigorous, weather-, drought-, and disease-resistant overstory; and grow trees for the next harvest. It will promote crown expansion, height growth, and diameter growth in the released stems. It should also focus on thinning from below for trees that are declining or poor quality co-dominants or sub-dominants. Crop tree release can be applied largely to maintain canopy closure and thereby reduce the likelihood of an increase in abundance of invasive plant species due to timber harvesting. This practice is a well-established silvicultural technique and often used on state lands. It has not been well tested for combating problematic invasive species. This will increase stand diversity and resiliency and minimize impacts of exotic plant species.

2. Use group selection (address Goals 6 and 8)

Use group selection to create gaps of sufficient size to allow regeneration to overtop invasive species before canopy closure and create additional age classes. Canopy gaps should be about two tree heights across or greater, at least 0.5 acres and up to 2 acres (VT Fish and Wildlife Department 2012). Proposed gaps should be treated for invasive species before and, if necessary, after logging (VT Fish and Wildlife Department 2012). This practice is a well-established silvicultural technique and often used on state lands, though not routinely used to combat problematic invasive species.

3. Treat unacceptable growing stock (UGS) (addresses Goal 8)

Most UGS should be harvested to eliminate cull or other low-vigor trees and retain healthy stems. This will improve species- and stand-level disease resistance by reducing stems vulnerable to disease, pests, drought, or storm damage. Operations should continue to follow already existing agency practices that aim for retaining about 20% of UGS to provide wildlife trees and future down woody material. Standing dead trees should be retained where possible as wildlife trees and for future down woody debris (Bennett 2010). These practices are already applied to state lands and to varying degrees on private lands.

4. Reduce eastern hemlock densities (addresses Goals 2, 3, and 6)

Eastern hemlock is highly susceptible to HWA, which has begun to invade the southern Vermont region. By eliminating over-mature and other low-quality hemlock trees (e.g., crown ratios $\geq 30\%$) and trees on shallow soils (Smith et al. 1997), the remaining hemlock trees are more likely to resist HWA and other pest species. Moreover, if HWA causes very high mortality, then increasing overstory diversity will prepare the stand to be able to recover more rapidly. Hemlock trees with crown ratios $> 30\%$ that are suppressed should be considered for release as the vigor of these trees should increase following release (Smith et al. 1997) and may become more resistant to HWA (Fajvan 2008). This practice could be used on state and private lands where eastern hemlock is dominant and risk for HWA is high in the next 15 to 25 years.

5. Apply soil BMPs (addresses Goals 1 and 2)

Carefully select time of year for harvest to control soil erosion, spread of exotic species, and create a seedbed for timber species. A winter harvest may reduce the likelihood of an expansion of invasive plant species by minimizing soil disturbance, which creates a seed bed for invasive plant species, and provides some protection to rare plant species

in the stand (VT TNC 2012). This practice is already used on state lands where invasive species are a problem. However, to create a seedbed for regenerating species that requires soil scarification for seedling establishment (e.g., yellow birch), a summer harvest will be necessary.

Potential Monitoring Practices

The goal of these monitoring practices is to collect information to support adaptive management actions that respond to changed forest conditions (see Table 1).

1. Monitor deer impacts with exclosures (addresses Goals 6 and 8)

Deer exclosures can be used to both assess and verify deer impacts and protect regeneration in harvest areas. Ideally three or more small exclosures should be used to determine the statistical significance of the impact of deer exclosures. Areas for placing the deer exclosures to be paired with non-exclosure sampling sites should be identified and sampled before harvesting (Year 0) and in Year 1, Year 5, Year 10, and Year 15 following the harvest. The following variables are recommended for monitoring: (1) stem counts of tree seedlings and saplings by species and height class, (2) stem counts of invasive plant species by species and height class, and (3) percent cover of a few herbaceous plant species that are sensitive to deer browsing (e.g., *Trillium* spp., *Maianthemum canadense*, *Medeola virginiana*) and fern spp. (VT Fish and Wildlife Department 2012). Depending on the size of exclosure, only a small portion of the exclosure would need to be sampled. This strategy emerged from discussions with the Stewardship Team. This strategy could occasionally be applied to state lands but would be too expensive to widely apply. These variables should indicate whether deer are having a significant impact on the understory plant communities by Year 10.

2. Monitor harvest gaps for regeneration and invasive species (addresses Goals 6 and 8)

Harvest gaps should be field checked one to two years after harvest to determine whether invasive plant species are suppressing the growth of regeneration (VT TNC 2012). Field checks by Year 2 should be able to identify whether control of exotic plant species in harvest gaps is necessary in order to maintain regeneration.

Table 1. Summary of potential monitoring and management actions related to climate change.

Attribute for Management	Potential Monitoring Action	When?	Estimated Monitoring Cost ¹	Potential Management Action	When?	Estimated Management Cost ¹
Deer Damage	Assess deer impacts on regeneration	Year 0, 1, 5, 10, & 15	\$750 ²	Install enclosures to protect patches of regeneration in harvest gaps	Year 1	\$2000-\$8000 (four enclosures; \$500-\$2000/33'x33'enclosures) ⁴
Invasive Plant Species	Assess regeneration and invasive plant species impacts on regeneration	Year 0, 1, 5, 10 & 15	\$750 ²	Apply herbicides using backpack sprayers to control exotic plant species	Year 2	\$35-\$245/acre ⁵
HWA	Assess impacts of HWA	Year 7 and 15	\$600	Consider pre-salvage and salvage operations	Year 15	No additional cost ³
Timber	Assess timber and regeneration	Year 15	No additional cost	Timber harvesting	Year 15	No additional cost

¹ Estimated costs include total cost of materials and labor over 15 years.
² Deer damage and invasive plant species monitoring would occur in combined visits, cost \$300/half-day visit.
³It is recommended that pre-salvage and salvage operations occur only as a part of routine timber harvest operations.
⁴ After Nicholas et al. 2007.
⁵ After MA Executive Office of Energy and Environmental Affairs. 2013, Kochenderfer et al. 2012.

3. Monitor for HWA (addresses Goals 6 and 8)

Eastern hemlock should be monitored for HWA every two to three years following Costa and Onken (2006) or similar protocols and for decline (percentage of healthy trees [$> 30\%$ crown ratio], declining trees [$\leq 30\%$ crown ratio], and dead trees). This strategy should be applied in areas at risk to HWA where eastern hemlock is dominant. Field surveys can be used to determine whether pre-salvage or salvage operations should be considered to promote forest regeneration and control fuel loads.

4. Review the site in 15 years for harvesting timber (addresses Goal 8)

The site should be evaluated in 15 to 20 years to determine whether a subsequent harvest consistent with land objectives at that time is possible and warranted. This should include an assessment of merchantable timber, regeneration, deer, exotic species, and pest species threats. This is a routine strategy for WMA stands with timber harvesting.

Potential Management Practices

The goal of these management actions is to adapt to three possible indirect impacts of climate change: increased deer populations due to milder winters, more invasive plants due to a more favorable growing season, and HWA due to milder winters (see Table 1).

1. Deer enclosures (addresses Goal 6 and 8)

Deer enclosures can be used to protect regeneration in harvest areas. Ideally three or more small enclosures could be used to protect seedlings and saplings in large harvest gaps. Enclosures should protect a diversity of regenerating tree species but should include red oak and other southern, drought tolerant species. This strategy could occasionally be applied to state lands but would be too expensive to apply widely. Alternatively more, small temporary enclosures could be applied (10' x 10').

2. Invasive plant species control in harvest gaps (addresses Goals 6 and 8)

Harvest gaps should be field checked one to two years after harvest to determine whether invasive plant species are suppressing the growth of regeneration (VT TNC 2012). If necessary, management staff should consider the application of herbicides or manual removal to reduce the impacts of invasive plant species. The staff had already planned to apply this strategy. This is a strategy that should be considered for all state and private lands that have been harvested where invasive plant species are a problem. Field checks by Year 2 should be able to identify whether control of exotic plant species in harvest gaps is necessary in order to maintain regeneration.

3. Salvage or pre-salvage of hemlock

Salvage or pre-salvage of hemlock should be considered in conjunction with routine planned harvests. Hemlock

surveys should be conducted in conjunction with pre-harvest timber surveys in Year 15 to determine whether salvage or pre-salvage of hemlock is necessary. Unsalvageable hemlock will contribute to the down woody material.

References

- Allen, C., A. Macalady, H. Chenchouni, D. Bachelet, N. McDowell, M. Vennetier, and N. Cobb. 2010. A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest Ecology and Management* 259: 660-684.
- Auclair, A., W. Heilman, and B. Brinkman. 2010. Predicting forest dieback in Maine, USA: a simple model based on soil frost and drought. *Can. J. For. Res.* 40: 687-702.
- Bennett, K. (editor). 2010. *Good Forestry in the Granite State: Recommended Voluntary Forest Management Practices for New Hampshire* (second edition). University of NH Cooperative Extension, Durham, NH.
- Burns, Russell M., and Barbara H. Honkala, tech. coords. 1990. *Silvics of North America: 1. Conifers; 2. Hardwoods*. USDAD, Agriculture Handbook 654. U.S. Department of Agriculture, Forest Service, Washington, DC. vol.2, 877 p.
- Butler, S., J. Melillo, J. Johnson, J. Mohan, P. Steudler, H. Lux, E. Burrows, R. Smith, C. Vario, L. Scott, T. Hill, N. Aponte, and F. Bowles. 2012. Soil warming alters nitrogen cycling in a New England forest: implications for ecosystem function and structure. *Oecologia* 168: 819-828.
- Costa, S. and B. Onken. 2006. Standardizing sampling for detection and monitoring of hemlock woolly adelgid in eastern hemlock forests. FHTET-20 06-16. Morgantown, WV: USDA, Forest Service, Forest Health Technology Enterprise Team, 11 pp.
- Dale, V., L. Joyce, S. McNulty, R. Neilson, M. Ayres, M. Flannigan, P. Hanson, L. Irland, A. Lugo, C. Peterson, D. Simberloff, F. Swanson, B. Stocks, and B. Wotton. 2001. Climate change can affect forests by altering the frequency, intensity, duration, and timing of fire, drought, introduced species, insect and pathogen outbreaks, hurricanes, windstorms, ice storms, or landslides. *Bioscience* 51:723-734.
- Drever, C., Y. Bergeron, M. Drever, M. Flannigan, T. Logan, and C. Messier. 2009. Effects of climate on occurrence and size of large fires in a northern hardwood landscape: historical trends, future predictions, and implications for climate change in Témiscamingue, Québec. *Applied Vegetation Science* 12:261-272.
- Dukes, J., and H. Mooney. 1999. Does global change increase the success of biological invaders? *Trends in Ecology and Evolution* 14: 135-139.
- Dukes, J., J. Pontius, D. Orwig, J. Garnas, V. Rodgers, N. Brazee, B. Cooke, K. Theoharides, E. Stange, R. Harrington, J. Ehrenfeld, J. Gurevitch, M. Lerda, K. Stinson, R. Wick, and M. Ayres. 2009. Response of insect pests, pathogens, and invasive plant species to climate change in the forests of northeastern North America: What can we predict? *Canadian Journal of Forest Research* 39:231-248.
- Fajvan, M. 2008. The role of silvicultural thinning in eastern forests threatened by hemlock woolly adelgid. In: Deal, R., ed. *Integrated restoration of forested ecosystems to achieve multi-resource benefits*. Proc. 2007 national silviculture workshop. Gen. Tech. Rep. 733. Portland, OR. USDA, Forest Service, Pacific Northwest Research Station: 247-256.
- Flannigan, M., M. Krawchuk, W. de Groot, B. Wotton and L. Gowman. 2009. Implications of changing climate for global wildland fire. *International Journal of Wildland Fire*. 18: 483-507.
- Frumhoff, P., J. McCarthy, J. Melillo, S. Moser, and D. Wuebbles. 2007. *Confronting Climate Change in the U.S. Northeast: Science, Impacts, and Solutions*. Synthesis report of the Northeast Climate Impacts Assessment. Cambridge, MA: Union of Concerned Scientists.
- Gunn, J., J. Hagan, and A. Whitman. 2009. *Forestry Adaptation and Mitigation in a Changing Climate: A forest resource manager's guide for the northeastern United States* (Report NCI-2009-1). Brunswick, ME: Manomet Center for Conservation Sciences.
- Hanson, P. and J. Weltzin. 2000. Drought disturbance from climate change: response of United States forests. *Science of the Total Environment* 262: 205-220.
- Hayhoe, K, C. Wake, T. Huntington, L. Luo, M. Schwartz, J. Sheffield, E. Wood, B. Anderson, I. Bradbury, T. DeGaetano, and D. Wolfe, D. 2007. Past and future changes in climate and hydrological indicators in the U.S. Northeast. *Climate Dynamics* 28: 381-407.
- Iverson, L.R., A.M. Prasad, and S. Matthews. 2008a. Modelling potential climate impacts on trees of the northeastern United States. *Mitigation and Adaptation Strategies for Global Change* 13:487-516.
- Iverson, L.R., A.M. Prasad, and M.W. Schwartz. 2005. Predicting potential changes in suitable habitat and distribution by 2100 for tree species of the eastern United States. *Journal of Agricultural Meteorology* 61:29-37.
- Knight, T., J. Dunn, L. Smith, J. Davis, and S. Kalisz. 2009. Deer facilitate invasive plant success in Pennsylvania forest understory. *Natural Areas Journal* 29: 110-116.
- Kochenderfer, J., J. Kochenderfer, and G. Miller. 2012. *Manual herbicide application methods for managing vegetation in Appalachian hardwood forests*. Gen. Tech. Rep. NRS 96. Newtown Square, PA. USDA, Forest Service, Northern Research Station. 59 p.
- MA Executive Office of Energy and Environmental Affairs. 2013. *LIP Habitat Management Practices Rates* (Based on NRCS WHIP Rates for 2012). MA Executive Office of Energy and Environmental Affairs, Boston, MA.
- Maciejowski, J., L. Thornton, S. Darling, D. Blodgett, G. Salmon, L. Richardson, M. Mayer, and C. MacKenzie.

2011. The Narrows Wildlife Management Area: Long Range Management Plan. VT ANR, Fish and Wildlife Department, Department of Forests, Parks & Recreation, Rutland North District Stewardship Team. Rutland, VT.
- Mohan, J., R. Cox, and L. Iverson. 2009. Composition and carbon dynamics of forests in northeastern North America in a future, warmer world. *Canadian Journal of Forest Research*. 39: 213-230.
- Nicholas, C., J. Wambaugh, T. Ladner, and J. Benner. 2007. Deer enclosure Guidelines. Bureau of Forestry, PA Department of Conservation & Nat. Res., Harrisburg, PA.
- Ollinger, S., C. Goodale, K. Hayhoe, and J. Jenkins. 2008. Potential effects of climate change and rising CO₂ on ecosystem processes in northeastern U.S. forests. *Mitigation and Adaptation Strategies for Global Change* 13: 467–485.
- Paradis A, J. Elkinton, K. Hayhoe, and I. Buonaccorsi. 2008. Role of winter temperature and climate change on the survival and future range expansion of the hemlock woolly adelgid (*Adelges tsugae*) in eastern North America. *Mitigation Adapt. Strateg. Global Change* 13:541-554.
- Prasad, A., L. Iverson., S. Matthews, and M. Peters. 2007-ongoing. A Climate Change Atlas for 134 Forest Tree Species of the Eastern United States [database]. <http://www.nrs.fs.fed.us/atlas/tree>, Northern Research Station, USDA Forest Service, Delaware, OH.
- Smith, D., B. Larson, M. Kelty, and P. Ashton. 1997. The practice of silviculture: Applied forest ecology. Wiley, New York. 537 p.
- Rustad, L.; J. Campbell; J. Dukes; T. Huntington; K. Lambert, J. Mohan, and N. Rodenhouse. 2012. Changing climate, changing forests: The impacts of climate change on forests of the northeastern United States and eastern Canada. Gen. Tech. Rep. NRS-99. Newtown Square, PA: USDA. Forest Service, Northern Research Station. 48 p.
- Tang, G. and B. Beckage. 2010. Projecting the distribution of forests in New England in response to climate change. *Diversity and Distributions* 16:144-158.
- USDA NRCS 2013. Soil Web. USDA NRCS, University of California, Davis, CA.
- VT Department of Forests, Parks, and Recreation. 2012. Vermont Invasive Forest Pest Update: Hemlock Woolly Adelgid *Adelges tsugae*. VT Department of Forests, Parks, and Recreation, Montpelier, VT.
- VT Fish and Wildlife Department. 2012. Deer Doing Damage to Land Managed for Production of Marketable Forest Products. Vermont Fish & Wildlife Department Working Group Report to Legislature - February 2012. Vermont Fish and Wildlife Department, Montpelier, VT.
- VT TNC. 2012. Best Management Practices for the Prevention and Treatment of Terrestrial Invasive Plants in Vermont Woodlands. TNC, Montpelier, VT.
- Wilkerson, E., H. Galbraith, A. Whitman, and S. Balch. 2011. Allen-Whitney Memorial Forest: Climate Change Adaptation Plan. Manomet Center for Conservation Sciences, Brunswick, ME.
- Zhu, K., C.W. Woodall, and J.C. Clark. 2012. Failure to migrate: lack of tree range expansion in response to climate change. *Global Change Biology* 18: 1042–1052.
- Ziska, L., and J. Dukes. 2011. *Weed Biology and Climate Change*. Oxford, UK: Wiley-Blackwell.

Acknowledgements

This report was greatly improved through review and comments from Sandy Wilmot, VT ANR, and Julie Beane and Ethel Wilkerson at Manomet. This report is partially supported by a grant from the USDA Forest Service to VT ANR and by a grant recommended by the USDA Forest Service, National Urban and Community Forestry Advisory Council in 2010.

Appendix A: Landowner Goals

The management goals within The Narrows WMA Long-range Management Plan (Maciejowski et al. 2011) are to:

1. Protect and enhance rare, threatened and endangered species and their habitat.
2. Maintain or enhance the quality of natural community conditions.
3. Protect and enhance wildlife habitat through management of all seral stages, creation of early successional growth, improvement of deer wintering areas, and protection of unique habitat.
4. Enhance opportunities for dispersed non-motorized activities for wildlife-based recreation, particularly hunting, trapping and wildlife viewing.
5. Protect and improve public access.
6. Demonstrate exemplary wildlife management practices so that practices applied here may find broader application on private lands.
7. Protect historic significance of the property, including known and suspected sites.
8. Provide sustainable, periodic timber harvesting in appropriate areas to promote wildlife habitat and forest productivity.